

Dietary fiber and reduced cough with phlegm: a cohort study in Singapore

Lesley M. Butler¹, Woon-Puay Koh², Hin-Peng Lee³, Mimi C. Yu², Stephanie J. London¹

Affiliations of authors: ¹Epidemiology Branch, National Institute of Environmental Health Sciences, Department of Health and Human Services, National Institutes of Health, Research Triangle Park, North Carolina; ²Department of Preventive Medicine, University of Southern California Keck School of Medicine, Los Angeles, California; ³National University of Singapore, Singapore.

Correspondence to: Dr. Lesley Butler, University of California, Department of Epidemiology and Preventive Medicine, One Shields Avenue, Davis, CA 95616, telephone: (530) 752-3898, fax: (530) 754-5621, e-mail: lmbutler@ucdavis.edu.

Reprint requests to: Dr. Stephanie London, NIEHS, PO Box 12233, MD A3-05, Research Triangle Park, NC, 27709, london2@niehs.nih.gov.

Running title: Fiber and reduced cough with phlegm. Subject code: 50, Chronic obstructive pulmonary disease (COPD): epidemiology. This article has an online supplement, which is accessible from this issue's table of contents online at www.atsjournals.org.

Word count: Introduction = 544; Methods = 790; Results = 917; Discussion = 1,613.

ABSTRACT

Smoking is the major risk factor for chronic respiratory symptoms, but dietary factors may also play a role. Most studies of diet and lung disease have been cross-sectional and conducted in populations with a Western-style diet. We analyzed the relation between dietary intake at baseline and new onset of cough with phlegm in a population-based cohort of 63,257 middle-aged Chinese men and women initiated in Singapore between 1993 and 1998. Beginning in 1999, we ascertained respiratory symptoms by telephone interview and have identified 571 incident cases of cough with phlegm among the 49,140 cohort members with completed follow-up. Non-starch polysaccharides (NSP), a major component of dietary fiber, total fruit and soy isoflavones had the strongest associations. Odds ratios comparing highest to lowest quartiles after adjustment for age, gender, dialect group, total energy intake and smoking were 0.61 (95% CI:0.47, 0.78, *P* for trend <0.001) for NSP, 0.67 (95% CI: 0.52, 0.87, *P* for trend=0.006) for fruit and 0.67 (95% CI:0.53, 0.86, *P* for trend=0.001) for soy isoflavones. These data suggest that a diet high in fiber from fruit and, possibly, soyfoods may reduce the incidence of chronic respiratory symptoms. Associated nutrients, such as flavonoids, may contribute to this association.

Keywords: dietary fiber, soy, fruit, chronic bronchitis, COPD

Word count: 199

Chronic respiratory symptoms are a major source of morbidity in adults (1). Cough and phlegm are frequently associated with chronic obstructive pulmonary disease (COPD) which may be caused by oxidative stress-mediated inflammation and tissue damage in the lung (2). Oxidants can cause direct damage by inactivating antiproteases or mediating other processes that promote the development of chronic lung damage (3). Cigarette smoke is a major source of exogenous oxidant exposure (4), and the major cause of chronic respiratory symptoms (5). However, susceptibility to respiratory morbidity varies among smokers and chronic cough and phlegm production and other respiratory-related morbidity occur in non-smokers (6). This suggests that additional factors, including other environmental exposures, genetics and diet may play a role. There is evidence that dietary nutrients, modulate oxidative stress-induced lung damage among both smokers and non-smokers (7-10).

Fruits and vegetables are the major food sources of antioxidants that may protect the lung from oxidative stress (11). Fruit intake and, to a lesser extent, vegetable intake have been associated with higher lung function and reduced symptoms of cough with phlegm (12,13). Cross-sectional studies support an association between vitamin C and lung function (14-18). However, the epidemiologic evidence is stronger for fruit intake than for individual fruit-related nutrients such as vitamin C and carotenoids (12,13), suggesting that other nutrients associated with fruit may be more relevant in protecting the lung from oxidative stressors.

In addition to vitamin C and carotenoids, fruit and vegetables contain flavonoids (19,20) and fiber. Flavonoids have free-radical scavenging properties that may influence lung disease (21,22), but have been rarely evaluated in prospective studies (11). Some epidemiologic evidence suggests that flavonoid-containing fruits, such as apples and pears, are associated with

improved lung function (23) and reduced non-specific respiratory disease (24). Reduced COPD symptoms was also reported among individuals with greater intake of the flavonoids known as catechins (25). In addition to fruit and vegetables, flavonoids are found in soyfoods (26,27), tea (28) and wine (29). Soyfoods are a major source of flavonoids in the Asian diet. Although experimental evidence suggests that genistein, a flavonoid in soyfoods, acts as an anti-inflammatory agent in the lung (30-32), the association between soyfoods and symptoms of chronic lung disease has not been investigated in an epidemiologic study. Dietary fiber is thought to contribute to the beneficial health effects of fruits and vegetables (33,34), but has also not been studied in relation to lung disease.

Prospective data are optimal to evaluate dietary relations with chronic lung disease and symptoms. The few prospective data that are available have limited detail about fruit and fiber intake. In addition, the available prospective data come exclusively from populations with Western-style diets (24,35-38). Studies in Asian populations may help to identify relations between lung disease and dietary factors, such as soy, which are uncommon elements of Western diets.

In a population-based cohort of Singaporeans of Chinese ethnicity, we assessed whether intake of fruit, vegetables, their associated nutrients, and other sources of flavonoids were associated with decreased risk of developing cough with phlegm. We used a dietary questionnaire, validated in our population, with extensive detail on the relevant foods (39). Some of the results of these analyses have been previously reported in the form of a poster presentation at the American Thoracic Society 2003 Conference (40).

METHODS

Study Population

The design of the Singapore Chinese Health Study has been described (41). Briefly, the cohort was drawn from men and women, aged 45 to 74 years, who were permanent residents or citizens of Singapore and resided in government-built housing estates (86% of the Singapore population reside in such facilities). We restricted the study to individuals belonging to the two major dialect groups of Chinese in Singapore: the Hokkiens and the Cantonese. Between April 1993 and December 1998, 63,257 individuals (~85% of those eligible) were enrolled. Enrollment in the cohort entailed completing a baseline in-person interview in the participants' home. The questionnaire elicited information on diet, demographics, current physical activity, reproductive history (women only), occupational exposure and medical history. Beginning in 1999 and ongoing, study participants responded to a single telephone interview including standardized questions on history of asthma and cough with phlegm production (42) (see online supplement for questionnaire items). The Institutional Review Boards at the University of Southern California, the National University of Singapore, and the National Institute of Environmental Health Sciences have approved this study.

As of December 31, 2002, there were 5,846 deaths in the overall cohort, with 4,871 of these deaths occurring before we were able to collect follow-up questionnaire data on respiratory outcomes. We analyzed data from the 49,140 individuals with completed telephone interviews as of August 3, 2003, with an average of 5.4 years of follow-up among cases (range=2.4 to 8.4; interquartile range = 4.4 to 6.4), and 5.2 years of follow-up among non-cases (range=1.2 to 9.2; interquartile range = 3.2 to 6.2). These 49,140 individuals represent 84.2% of the cohort available for respiratory outcome follow-up. (Demographic, smoking and dietary differences

between those with and without respiratory outcome data are presented in the online-supplement in Table E1). Incident cases were subjects who reported that they have cough with phlegm on arising or during the rest of the day, initiated after the baseline interview (N=571). In a subset analysis, cases were further defined by having incident cough with phlegm for three or more months of the year (N=359). We classified cough with phlegm production as incident with respect to baseline, if the difference in years between the age at the follow-up interview, and the duration of concurrent cough with phlegm, was greater than the age at the baseline interview. Non-cases (N=44,068) were cohort members without cough or phlegm. The remaining 4,498 individuals had either incident cough only (N=512), incident phlegm only (N=1,271), prevalent cough with phlegm (N=668) or discordant incident/prevalent combinations of cough and/or phlegm (N=2047). Analyses of the 4, 498 subjects with these secondary outcomes can be found in the online supplement (Tables E2-E6).

Dietary Assessment

At baseline, a 165-item quantitative food frequency questionnaire, developed for, and validated in, this population, along with color photographs to represent portion sizes, was used to assess usual diet over the past year (39). The questionnaire included 16 items on individual fruits and fruit juices, 24 items covering 34 specific vegetables, and seven soyfood items (39). Total soyfood intake was measured as the summation of all soyfoods expressed in units of plain tofu equivalent, to account for the different percentage of water in each soyfood, as previously described (43). In a validation substudy, frequency of overall soy intake had a statistically significant dose-dependent association with urinary total isoflavones (daidzein, genistein and glycitein) (44). The questionnaire also included items about black tea (Chinese red and ceylon tea) and green tea (jasmine and woolong).

Statistical analysis

We examined diet at baseline, in relation to the outcome of incident cough with phlegm reported on the follow-up questionnaire. Adjusted odds ratios and 95 percent confidence intervals were calculated from unconditional logistic regression models (version 8.1; SAS Institute, Cary, NC). All nutrient data were adjusted for total energy intake using the residual method (45). We present intake of non-starch polysaccharides, a major component of dietary fiber, instead of total dietary fiber to facilitate comparison across study populations (46,47). We adjusted for age, sex and dialect group because these were associated with the outcome although they did not necessarily appreciably alter the dietary associations. As expected, smoking was a strong risk factor for cough with phlegm in our study, therefore we sought to carefully control for potential confounding by smoking by including terms for smoking status (never, current, past), age at starting to smoke and amount smoked in adjusted models with dietary factors. Individuals were categorized as never smokers if they had never smoked at least one cigarette a day for one year or longer. Length of follow-up, defined as the interval from baseline to follow-up interview (48), was not included in the adjusted models, because it was not associated with the outcome and did not appreciably alter the dietary associations.

RESULTS

Cohort members with incident cough with phlegm were more likely to be male, older, and members of the Hokkien dialect group, compared to non-cases (Table 1). Cases, regardless of symptom duration, were more likely to be current smokers, to have started smoking before age 20, and were heavier smokers (Table 1). The mean total energy intake was slightly higher among all cases, compared to non-cases [1621.9 kcal, standard deviation (SD) = 572.2 among cases, and

1554.9, SD = 556.1 among non-cases], although it was not associated with the outcome after adjustment for smoking. Characteristics that did not differ substantially by case status included body mass index (mean = 22.9 kg/m², SD = 3.5 among cases, and mean = 23.2 kg/m², SD = 3.3 among non-cases), education level (22.6% cases and 26.2% non-cases had no formal education), and physical activity (7.8% cases and 8.4% non-cases with 4 hours per week of moderate activity).

Table 2 shows the associations between individual food and nutrient items and incident cough with phlegm adjusted, as were all subsequent models, for age, gender, dialect group and smoking (never, past, or current smoking status, age at starting to smoke and amount smoked). We did not include a separate term for years since quitting smoking or change in smoking status from baseline to follow-up in the adjusted models, because neither factor was not an important predictor of the outcome. For years since quitting, this was consistent with the approximately equal proportion of past smokers among the cases and non-cases. The associations did not substantially differ for any of the foods and nutrients when cases were restricted to those with cough plus phlegm lasting for three or more months out of the year (Table 2), therefore individual results reported in the text refer to all 571 incident cases. The associations that we observed for the incident cough with phlegm category were generally not seen for those with incident cough only, incident phlegm only, or for the miscellaneous category (see online supplement Tables E3-E5). The prevalent cough with phlegm category generally had findings similar to the incident category (see online supplement Tables E3-E5).

Daily intake of non-starch polysaccharides was inversely associated with incident cough plus phlegm [odds ratio (OR) = 0.61, 95% confidence interval (CI): 0.47, 0.78 comparing highest to lowest quartiles; *P* for trend <0.001] (Table 2). Total fruit, all grain products and total

vegetables were the greatest food sources of non-starch polysaccharides, with each contributing 34.3%, 24.6% and 22.2% of intake respectively. Total fruit was also inversely associated with cough plus phlegm, but all grain products and total vegetable consumption were not (Table 2).

When we looked at specific antioxidant nutrients related to fruit intake, such as vitamin C, we found trends similar to those observed for fruit (Table 2). However, we found no association for carotenoids, including total carotenoids, beta-carotene (Table 2), or other individual carotenoids, such as alpha-carotene, beta-cryptoxanthin, lycopene or lutein (data not shown). We also found no association for vitamin E (Table 2), magnesium or vitamin B-6 (data not shown). Given the low frequency of vitamin or mineral supplement use (8.4% in cases and 8.2% in non-cases), odds ratios for individual nutrients did not change when supplement users were removed from the models (data not shown).

Given the association with total fruit consumption, we examined associations with individual fruits. We categorized intake by tertiles rather than quartiles, because the frequency of consumption of any individual fruit was, as expected, lower than for total fruit. We observed the strongest inverse dose-response associations for apples, pears, and grapes (Table 3).

Soyfoods made the next greatest contribution to non-starch polysaccharide intake after fruit, grain products and vegetables. Total soyfood and soy isoflavone intake had modest inverse associations with cough plus phlegm (Table 4). Comparing the fourth to the first quartiles, odds ratios were virtually identical for the individual isoflavones genistein (OR = 0.68; 95% CI: 0.53, 0.87), daidzein (OR = 0.67; 95% CI: 0.53, 0.86,) and glycitein (OR = 0.68; 95% CI: 0.54, 0.87).

We examined whether associations between cough with phlegm production and individual foods and nutrients were altered by adjustment for total non-starch polysaccharides (Table 5). Odds ratios for all dietary factors were attenuated after adjustment for non-starch

polysaccharides; odds ratios for all grain products remained near the null. The inverse association between apples (OR = 0.79; 95% CI: 0.64, 0.98, comparing third to first tertile) and grapes (OR = 0.79; 95% CI: 0.62, 0.99, comparing third to first tertile) persisted, after adjustment for non-starch polysaccharides. With adjustment for the other foods and nutrients in Tables 2 through 4, the odds ratios for non-starch polysaccharides, comparing extreme quartiles, ranged from 0.49 to 0.67 and remained statistically significant.

There was no appreciable difference in association between incident cough with phlegm and non-starch polysaccharides by strata of age (< / ≥ median of 55 years), gender, dialect group (Hokkien / Cantonese) or body mass index (< / ≥ median of 23.1 kg/m²) (data not shown). Although we observed a stronger association for non-starch polysaccharide intake among ever smokers (OR = 0.45; 95% CI:0.29, 0.70, comparing fourth to first quartile), an inverse association was still present among never smokers (OR = 0.70; 95% CI:0.50, 0.97, comparing fourth to first quartile). There were no changes in the odds ratios for non-starch polysaccharide intake when individuals who self-reported incident or prevalent asthma (N=68 cases and 1719 non-cases) were removed (data not shown). Frequency of asthma by incident cough with phlegm case status is shown in the online supplement (Table E6).

DISCUSSION

These prospective data support the role of diet in the etiology of cough with phlegm production. In particular, non-starch polysaccharides, a major component of dietary fiber, and certain non-citrus fruits (e.g. apples, grapes) were independently associated with reduced development of cough with phlegm. As in previous prospective studies of lung function (35) or chronic respiratory symptoms and disease (24,36), we found modest inverse associations for fruit and

vitamin C intake. However, these associations were eliminated after adjustment for non-starch polysaccharide intake. The earlier prospective studies (24,35,36) did not consider the role of fiber, therefore it is unclear whether the previously reported associations with fruit and vitamin C were independent of non-starch polysaccharide intake.

Fruit, vegetable and grain products were the major contributors of non-starch polysaccharide intake in this population. Incidence of cough with phlegm was reduced with high intake of fruit, but not vegetable or grain products. The lack of an inverse association with vegetable intake may be due to its limited variation in our population, where the interquartile range was 2.6-fold smaller for vegetable than for fruit intake. The limited variation would give us much less power to find an association with vegetables than with fruit.

It is also possible that fiber from fruit may have physiologic effects that are more beneficial in reducing development of cough with phlegm than fiber from vegetables or grain products. In addition, to the extent that fiber from whole grains is important, we might not see this in our cohort because white rice is the major grain in the Asian diet, rather than the whole grains found in some Western diets (49). Fiber is a complex dietary component (47,50) and may have beneficial effects on the lung such as reducing blood glucose concentrations (51-53), thereby reducing inflammation (54) and enhancing antioxidant processes (33,34). The physiologic properties of fiber from fruits versus vegetables and grains in relation to lung disease have not been studied. However, in support of our finding for respiratory symptoms, fiber from fruit had the greatest inverse association with risk of coronary heart disease in a recent pooled analysis, compared to other sources of dietary fiber (55).

Another hypothesis for why fruit, but not vegetable or grain intake, was inversely associated with cough plus phlegm is that components in fruit, in addition to fiber, such as

individual flavonoids, may be important in protection against chronic productive cough.

Flavonoids may protect the lung based on their antioxidant (56) and anti-inflammatory properties (22,57). Flavonoid antioxidant mechanisms may involve direct free-radical scavenging and inhibition of nitric-oxide synthase or xanthine oxidase (22,58). In addition, flavonoid anti-inflammatory mechanisms involve inhibitory effects *in vitro* on the cyclooxygenase and 5-lipoxygenase pathways of arachidonic acid metabolism (59,60), including inhibition of protein kinases (61).

Flavonoids are a diverse group of polyphenols (56,62). Although both fruits and vegetables are sources of flavonoids, particular classes of these compounds are more abundant in fruits (63,64). Predominant flavonoids in the diet include quercetin, a flavonol primarily found in apples, berries, tea and onions (65), and catechin, a flavanol primarily found in apples, pears, black grapes, tea and apricots (66). In the Asian diet, with high intake of tofu soy isoflavones are an important source of flavonoids (44,67). Food composition databases to assess flavonoid intake have recently been developed for the U.S. (63,65) and The Netherlands (64,68,69), but not for the diet of Asian countries. Thus we cannot evaluate total flavonoid intake in our study population at this time.

Our findings suggest that flavonoids may play a crucial role in the relation between fruit and soyfood intake and development of chronic cough with phlegm. We observed inverse associations between incident cough with phlegm and fruits high in flavonoids, such as apples, grapes and pears (65,66), and the associations with apples and grapes remained statistically significant after adjustment for total non-starch polysaccharide intake. An inverse association between flavonoids and lung disease was first suggested by Miedema, *et al* based on findings from a prospective study in which intake of apples and pears reduced the risk of chronic non-

specific lung disease; no association was found for vitamin C or beta-carotene (24). Two studies suggest that higher intake of apples and wine is associated with improved lung function (23,70). In a cross-sectional study, Tabak, *et al* found intake of three classes of flavonoids (catechins, flavonols and flavones) to be related to lung function, chronic cough, and breathlessness, but not with chronic phlegm (25). Solid fruit intake was associated with a decreased risk of all three symptoms (25).

Additional support for the importance of flavonoids comes from our finding of an inverse association with soyfoods and soy isoflavones. We are not aware of any previous epidemiologic data on soyfoods and non-malignant lung disease. A role for soy is plausible, because genistein, a soy isoflavone, acts *in vivo* as an anti-inflammatory agent by inhibiting protein tyrosine kinase (30) and nuclear factor-kappa B (31). Daily tea drinking, a source of flavonoids, had a weak inverse association with cough plus phlegm (data not shown). Wine, another source of flavonoids that may have beneficial effects on the lung (70,71), cannot be assessed in our population because only 0.8% drank at least one glass of (red or white) wine annually.

A limitation of observational studies is the potential for differential recall bias of usual diet by case status. With specific reference to our population, traditional Chinese medicine recommends that individuals with cough, wheeze and/or excess phlegm, avoid “Yin” foods, or foods with “cold” properties (72,73). Most fruits and vegetables are considered Yin (72). Although we prospectively assessed cough with phlegm, we considered the possibility that individuals may have avoided certain foods, which could cause a false protective association. However, after removing individuals who reported onset of symptoms within the first year after the baseline diet interview, we observed no differences in the odds ratios for total fruit or total non-starch polysaccharide intake with cough plus phlegm (data not shown). Further, if this bias

were operating, we would have expected to see an inverse association with vegetable intake, which we did not. We concluded that our findings could not be explained by avoidance of Yin foods due to early symptoms. This possible bias was not relevant to soy which is considered “neutral” or without Yin and Yang characteristics.

One possible concern is that the presence of cough with phlegm may be due to infectious causes. It should be noted that the period of symptom ascertainment (July 1999 – August 2003) preceded the recent severe acute respiratory syndrome outbreak. Further, previous diagnosis of tuberculosis was rare among our case group (2.3%, with average age at diagnosis of 37 years).

The objection may be raised that a diet high in fiber, fruit and soy might be a marker of a healthy lifestyle, including lower smoking (74,75). We carefully adjusted for smoking, although eliminating residual confounding is difficult to guarantee. However, we found a protective effect of non-starch polysaccharide intake in both smokers and nonsmokers suggesting that the protective effects of non-starch polysaccharides are not due to correlations with smoking behavior.

We had follow-up data on respiratory outcomes for 84.2% of the Singapore Chinese Health Study participants. We would expect that some differences would exist between those with and without respiratory outcome data, some of whom are deceased (see online supplement Table E1). However, for loss to follow-up to have created the inverse association we observed, we would have had to disproportionately lose either cases with higher fiber intake, or non-cases with low fiber intake. While we obviously lack respiratory data on those who have yet to complete the follow-up interview or died prior to completion, when we compare the dietary data at baseline among participants with and without follow-up, participants without outcome data have similar to slightly lower intakes for all the key nutrients (see online supplement Table E1).

Thus, it is not plausible that the lack of follow-up data in this cohort could be responsible for the observed inverse associations between diet and incident cough with phlegm.

The strengths of this study include its prospective design, which allows us to have greater confidence that associations we observe between diet and symptoms reflect effects on incidence rather than recall bias or dietary changes due to symptom development. We acknowledge that for symptoms of cough and phlegm, subjects will have some difficulty reporting the exact number of years they have had the symptoms and thus there will be some misclassification between the prevalent and incident categories. However, on average, individuals who we classified as "incident" are much more likely to have initiated symptoms after the baseline dietary interview compared to individuals we classified as "prevalent."

Other advantages of the study include high quality dietary data, specifically the use of a 165-item food frequency questionnaire, developed for, and validated in, this population, and extensive smoking and other lifestyle data to adequately adjust for potential confounding effects. A unique aspect of these analyses was the Asian population that enabled us to examine foods less common in a Western-style diet, such as soy. Vitamin supplement use was also low compared with U.S. populations, enabling us to better examine effects of food intake.

Preventing the initiation of smoking is still the most important primary prevention method for chronic respiratory symptoms and COPD in general, not only because it is the major risk factor for development of the conditions, but also because adverse effects on lung parenchyma remain long after cessation (76,77). However, our data from the Singapore Chinese Health Study provide evidence that promoting a diet high in sources of fiber and flavonoids, such as fruit and soy, may be an important contribution to primary prevention strategies for chronic respiratory symptoms, both in smokers and non-smokers.

ACKNOWLEDGMENTS

We thank Siew-Hong Low of the National University of Singapore for supervising the field work of the Singapore Chinese Health Study, Kazuko Arakawa of the University of Southern California for the development and management of the cohort study database, Marsha Shepherd of Westat, Inc. for assistance with programming, and Dr. David Dunson of the National Institute for Environmental Health Sciences for statistical advice. This research was supported by the U.S. National Institutes of Health (NCI RO1 CA80205 and NIEHS ZO1 ES43012).

REFERENCES

1. McSweeney, AJ, Grant, I, Heaton, RK, Adams, KM, Timms, RM. Life quality of patients with chronic obstructive pulmonary disease. *Arch Intern Med* 1982;142:473-478.
2. Heffner, JE, Repine, JE. Pulmonary strategies of antioxidant defense. *Am Rev Respir Dis* 1989;140:531-554.
3. Repine, JE, Bast, A, Lankhorst, I. Oxidative stress in chronic obstructive pulmonary disease. Oxidative Stress Study Group. *Am J Respir Crit Care Med* 1997;156:341-357.
4. Pryor, WA, Stone, K. Oxidants in cigarette smoke. Radicals, hydrogen peroxide, peroxyxynitrate, and peroxyxynitrite. *Ann N Y Acad Sci* 1993;686:12-27; discussion 27-18.
5. Barnes, PJ. Chronic obstructive pulmonary disease. *N Engl J Med* 2000;343:269-280.
6. Mannino, DM. COPD: epidemiology, prevalence, morbidity and mortality, and disease heterogeneity. *Chest* 2002;121:121S-126S.
7. Smit, HA. Chronic obstructive pulmonary disease, asthma and protective effects of food intake: from hypothesis to evidence? *Respir Res* 2001;2:261-264.
8. Burney, P. The origins of obstructive airways disease. A role for diet? *Am J Respir Crit Care Med* 1995;151:1292-1293.
9. MacNee, W. Oxidants/antioxidants and COPD. *Chest* 2000;117:303S-317S.
10. Altose, MD. Approaches to slowing the progression of COPD. *Curr Opin Pulm Med* 2003;9:125-130.
11. Schunemann, H, Freudenheim, J, Grant, B. Epidemiologic evidence linking antioxidant vitamins to pulmonary function and airway obstruction. *Epidemiol Rev* 2001;23:248-267.
12. Romieu, I, Trenga, C. Diet and obstructive lung diseases. *Epidemiol Rev* 2001;23:268-287.

13. Smit, HA, Grievink, L, Tabak, C. Dietary influences on chronic obstructive lung disease and asthma: a review of the epidemiological evidence. *Proc Nutr Soc* 1999;58:309-319.
14. Chen, R, Tunstall-Pedoe, H, Bolton-Smith, C, Hannah, MK, Morrison, C. Association of dietary antioxidants and waist circumference with pulmonary function and airway obstruction. *Am J Epidemiol* 2001;153:157-163.
15. Hu, G, Cassano, PA. Antioxidant nutrients and pulmonary function: the Third National Health and Nutrition Examination Survey (NHANES III). *Am J Epidemiol* 2000;151:975-981.
16. Grievink, L, Smit, HA, Ocke, MC, van 't Veer, P, Kromhout, D. Dietary intake of antioxidant (pro)-vitamins, respiratory symptoms and pulmonary function: the MORGEN study. *Thorax* 1998;53:166-171.
17. Britton, JR, Pavord, ID, Richards, KA, Knox, AJ, Wisniewski, AF, Lewis, SA, Tattersfield, AE, Weiss, ST. Dietary antioxidant vitamin intake and lung function in the general population. *Am J Respir Crit Care Med* 1995;151:1383-1387.
18. Hu, G, Zhang, X, Chen, J, Peto, R, Campbell, TC, Cassano, PA. Dietary vitamin C intake and lung function in rural China. *Am J Epidemiol* 1998;148:594-599.
19. Nielsen, SE, Freese, R, Kleemola, P, Mutanen, M. Flavonoids in human urine as biomarkers for intake of fruits and vegetables. *Cancer Epidemiol Biomarkers Prev* 2002;11:459-466.
20. Miesan, KH, Mohamed, S. Flavonoid (myricetin, quercetin, kaempferol, luteolin, and apigenin) content of edible tropical plants. *J Agric Food Chem* 2001;49:3106-3112.
21. Lu, FJ, Chu, LH, Gau, RJ. Free radical-scavenging properties of lignin. *Nutr Cancer* 1998;30:31-38.

22. Nijveldt, RJ, van Nood, E, van Hoorn, DE, Boelens, PG, van Norren, K, van Leeuwen, PA. Flavonoids: a review of probable mechanisms of action and potential applications. *Am J Clin Nutr* 2001;74:418-425.
23. Butland, BK, Fehily, AM, Elwood, PC. Diet, lung function, and lung function decline in a cohort of 2512 middle aged men. *Thorax* 2000;55:102-108.
24. Miedema, I, Feskens, EJ, Heederik, D, Kromhout, D. Dietary determinants of long-term incidence of chronic nonspecific lung diseases. The Zutphen Study. *Am J Epidemiol* 1993;138:37-45.
25. Tabak, C, Arts, IC, Smit, HA, Heederik, D, Kromhout, D. Chronic obstructive pulmonary disease and intake of catechins, flavonols, and flavones: the MORGEN Study. *Am J Respir Crit Care Med* 2001;164:61-64.
26. Franke, AA, Hankin, JH, Yu, MC, Maskarinec, G, Low, SH, Custer, LJ. Isoflavone levels in soy foods consumed by multiethnic populations in Singapore and Hawaii. *J Agric Food Chem* 1999;47:977-986.
27. Lampe, JW, Gustafson, DR, Hutchins, AM, Martini, MC, Li, S, Wahala, K, Grandits, GA, Potter, JD, Slavin, JL. Urinary isoflavonoid and lignan excretion on a Western diet: relation to soy, vegetable, and fruit intake. *Cancer Epidemiol Biomarkers Prev* 1999;8:699-707.
28. Balentine, DA, Wiseman, SA, Bouwens, LC. The chemistry of tea flavonoids. *Crit Rev Food Sci Nutr* 1997;37:693-704.
29. Scalbert, A, Williamson, G. Dietary intake and bioavailability of polyphenols. *J Nutr* 2000;130:2073S-2085S.

30. Duan, W, Kuo, IC, Selvarajan, S, Chua, KY, Bay, BH, Wong, WS. Anti-inflammatory Effects Of Genistein, A Tyrosine Kinase Inhibitor, On A Guinea Pig Model Of Asthma. *Am J Respir Crit Care Med* 2003;167:185-192.
31. Kang, JL, Lee, HW, Lee, HS, Pack, IS, Chong, Y, Castranova, V, Koh, Y. Genistein prevents nuclear factor-kappa B activation and acute lung injury induced by lipopolysaccharide. *Am J Respir Crit Care Med* 2001;164:2206-2212.
32. Akiyama, T, Ishida, J, Nakagawa, S, Ogawara, H, Watanabe, S, Itoh, N, Shibuya, M, Fukami, Y. Genistein, a specific inhibitor of tyrosine-specific protein kinases. *J Biol Chem* 1987;262:5592-5595.
33. Eastwood, MA. Interaction of dietary antioxidants in vivo: how fruit and vegetables prevent disease? *Qjm* 1999;92:527-530.
34. Larrauri, JA, Goni, I, Martin-Carron, N, Ruperez, P, Saura-Calixto, F. Measurement of health-promoting properties in fruit dietary fibers: antioxidant capacity, fermentability and glucose retardation index. *J Science Food Agric* 1996;71:515-519.
35. Carey, IM, Strachan, DP, Cook, DG. Effects of changes in fresh fruit consumption on ventilatory function in healthy British adults. *Am J Respir Crit Care Med* 1998;158:728-733.
36. Walda, IC, Tabak, C, Smit, HA, Rasanen, L, Fidanza, F, Menotti, A, Nissinen, A, Feskens, EJ, Kromhout, D. Diet and 20-year chronic obstructive pulmonary disease mortality in middle-aged men from three European countries. *Eur J Clin Nutr* 2002;56:638-643.
37. McKeever, TM, Scrivener, S, Broadfield, E, Jones, Z, Britton, J, Lewis, SA. Prospective study of diet and decline in lung function in a general population. *Am J Respir Crit Care Med* 2002;165:1299-1303.

38. Butland, BK, Strachan, DP, Anderson, HR. Fresh fruit intake and asthma symptoms in young British adults: confounding or effect modification by smoking? *Eur Respir J* 1999;13:744-750.
39. Hankin, JH, Stram, DO, Arakawa, K, Park, S, Low, SH, Lee, HP, Yu, MC. Singapore Chinese Health Study: development, validation, and calibration of the quantitative food frequency questionnaire. *Nutr Cancer* 2001;39:187-195.
40. Butler, L, Koh, WP, Lee, HP, Yu, MC, London, SJ. Foods high in soluble fiber may reduce risk of chronic cough and phlegm: the Singapore Chinese cohort study [abstract]. *Am J Respir Crit Care Med* 2003;167:A443.
41. Yuan, JM, Stram, DO, Arakawa, K, Lee, HP, Yu, MC. Dietary cryptoxanthin and reduced risk of lung cancer: the Singapore Chinese Health Study. *Cancer Epidemiol Biomarkers Prev* 2003;12:890-898.
42. Ferris, BG. Epidemiology Standardization Project (American Thoracic Society). *Am Rev Respir Dis* 1978;118:1-120.
43. Sun, CL, Yuan, JM, Arakawa, K, Low, SH, Lee, HP, Yu, MC. Dietary soy and increased risk of bladder cancer: the Singapore Chinese Health Study. *Cancer Epidemiol Biomarkers Prev* 2002;11:1674-1677.
44. Seow, A, Shi, CY, Franke, AA, Hankin, JH, Lee, HP, Yu, MC. Isoflavonoid levels in spot urine are associated with frequency of dietary soy intake in a population-based sample of middle-aged and older Chinese in Singapore. *Cancer Epidemiol Biomarkers Prev* 1998;7:135-140.
45. Willett, W, Stampfer, MJ. Total energy intake: implications for epidemiologic analyses. *Am J Epidemiol* 1986;124:17-27.

46. Goodlad, RA, Englyst, HN. Redefining dietary fibre: potentially a recipe for disaster. *Lancet* 2001;358:1833-1834.
47. Institute of Medicine. Dietary Reference Intakes: Proposed Definition of Dietary Fiber: Washington, D.C.: National Academy Press; 2001. 64 p.
48. Dinse, G, Lagakos, S. Regression analysis of tumour prevalence data. *Applied Statistics* 1983;32:236-248.
49. Chen, HL, Huang, YC. Fiber intake and food selection of the elderly in Taiwan. *Nutrition* 2003;19:332-336.
50. Lee, SC, Prosky, L. International survey on dietary fiber: definition, analysis, and reference materials. *J AOAC Int* 1995;78:22-36.
51. Walter, RE, Beiser, A, Givelber, RJ, O'Connor, GT, Gottlieb, DJ. Association between glycemic state and lung function: the Framingham Heart Study. *Am J Respir Crit Care Med* 2003;167:911-916.
52. Eliraz, A, Sternfeld, M, Madar, Z. Glycaemic and insulinaemic index of meals in stable obstructive pulmonary disease subjects. *Eur J Clin Nutr* 1994;48:522-524.
53. Davis, WA, Knuiaman, M, Kendall, P, Grange, V, Davis, TM. Glycemic Exposure Is Associated With Reduced Pulmonary Function in Type 2 Diabetes: The Fremantle Diabetes Study. *Diabetes Care* 2004;27:752-757.
54. Liu, S, Manson, JE, Buring, JE, Stampfer, MJ, Willett, WC, Ridker, PM. Relation between a diet with a high glycemic load and plasma concentrations of high-sensitivity C-reactive protein in middle-aged women. *Am J Clin Nutr* 2002;75:492-498.

55. Pereira, MA, O'Reilly, E, Augustsson, K, Fraser, GE, Goldbourt, U, Heitmann, BL, Hallmans, G, Knekt, P, Liu, S, Pietinen, P, *et al.* Dietary fiber and risk of coronary heart disease: a pooled analysis of cohort studies. *Arch Intern Med* 2004;164:370-376.
56. Pietta, PG. Flavonoids as antioxidants. *J Nat Prod* 2000;63:1035-1042.
57. Robak, J, Gryglewski, RJ. Bioactivity of flavonoids. *Pol J Pharmacol* 1996;48:555-564.
58. Bors, W, Saran, M. Radical scavenging by flavonoid antioxidants. *Free Radic Res Commun* 1987;2:289-294.
59. Laughton, MJ, Evans, PJ, Moroney, MA, Hoult, JR, Halliwell, B. Inhibition of mammalian 5-lipoxygenase and cyclo-oxygenase by flavonoids and phenolic dietary additives. Relationship to antioxidant activity and to iron ion-reducing ability. *Biochem Pharmacol* 1991;42:1673-1681.
60. Kim, HP, Mani, I, Iversen, L, Ziboh, VA. Effects of naturally-occurring flavonoids and biflavonoids on epidermal cyclooxygenase and lipoxygenase from guinea-pigs. *Prostaglandins Leukot Essent Fatty Acids* 1998;58:17-24.
61. Manthey, JA. Biological properties of flavonoids pertaining to inflammation. *Microcirculation* 2000;7:S29-34.
62. Heim, KE, Tagliaferro, AR, Bobilya, DJ. Flavonoid antioxidants: chemistry, metabolism and structure-activity relationships. *J Nutr Biochem* 2002;13:572-584.
63. Peterson, J, Dwyer, J. Flavonoids: dietary occurrence and biochemical activity. *Nutr Res* 1998;18:1995-2018.
64. Hertog, MG, Hollman, PCH, Katan, MB. Content of potentially anticarcinogenic flavonoids of 28 vegetables and 9 fruits commonly consumed in The Netherlands. *J Agric Food Chem* 1992;40:2379-2283.

65. Sampson, L, Rimm, E, Hollman, PC, de Vries, JH, Katan, MB. Flavonol and flavone intakes in US health professionals. *J Am Diet Assoc* 2002;102:1414-1420.
66. Arts, IC, Hollman, PC, Bueno De Mesquita, HB, Feskens, EJ, Kromhout, D. Dietary catechins and epithelial cancer incidence: the Zutphen elderly study. *Int J Cancer* 2001;92:298-302.
67. Chen, Z, Zheng, W, Custer, LJ, Dai, Q, Shu, XO, Jin, F, Franke, AA. Usual dietary consumption of soy foods and its correlation with the excretion rate of isoflavonoids in overnight urine samples among Chinese women in Shanghai. *Nutr Cancer* 1999;33:82-87.
68. Arts, IC, van de Putte, B, Hollman, PC. Catechin contents of foods commonly consumed in The Netherlands. 1. Fruits, vegetables, staple foods, and processed foods. *J Agric Food Chem* 2000;48:1746-1751.
69. Arts, IC, van De Putte, B, Hollman, PC. Catechin contents of foods commonly consumed in The Netherlands. 2. Tea, wine, fruit juices, and chocolate milk. *J Agric Food Chem* 2000;48:1752-1757.
70. Schunemann, HJ, Grant, BJ, Freudenheim, JL, Muti, P, McCann, SE, Kudalkar, D, Ram, M, Nochajski, T, Russell, M, Trevisan, M. Beverage specific alcohol intake in a population-based study: Evidence for a positive association between pulmonary function and wine intake. *BMC Pulm Med* 2002;2:3.
71. Shaheen, SO, Sterne, JA, Thompson, RL, Songhurst, CE, Margetts, BM, Burney, PG. Dietary antioxidants and asthma in adults: population-based case-control study. *Am J Respir Crit Care Med* 2001;164:1823-1828.

72. Weng, W, Chen, J. The eastern perspective on functional foods based on traditional Chinese medicine. *Nutr Rev* 1996;54:S11-16.
73. Connett, GJ, Lee, BW. Treating childhood asthma in Singapore: when West meets East. *Bmj* 1994;308:1282-1284.
74. Wynder, EL, Stellman, SD, Zang, EA. High fiber intake. Indicator of a healthy lifestyle. *Jama* 1996;275:486-487.
75. Margetts, BM, Jackson, AA. Interactions between people's diet and their smoking habits: the dietary and nutritional survey of British adults. *Bmj* 1993;307:1381-1384.
76. Rutgers, SR, Postma, DS, ten Hacken, NH, Kauffman, HF, van Der Mark, TW, Koeter, GH, Timens, W. Ongoing airway inflammation in patients with COPD who do not currently smoke. *Thorax* 2000;55:12-18.
77. Turato, G, Di Stefano, A, Maestrelli, P, Mapp, CE, Ruggieri, MP, Roggeri, A, Fabbri, LM, Saetta, M. Effect of smoking cessation on airway inflammation in chronic bronchitis. *Am J Respir Crit Care Med* 1995;152:1262-1267.

Table 1 Characteristics of the Singapore Chinese Health Study for subjects with and without incident cough with phlegm*

	Non-cases N=44,068	All cases of incident cough with phlegm N=571‡	Cases with symptoms for 3 months per year N=359‡
Sex (%)			
Men	40.3	58.3	58.5
Women	59.7	41.7	41.5
Age			
Mean years (standard deviation)	55.7 (7.7)	58.0 (7.9)	58.0 (8.1)
Dialect group (%)			
Hokkien	51.1	58.5	67.7
Cantonese	48.9	41.5	32.3
Smoking status (%)			
Never§	74.3†	49.9	49.3
Past	10.3	12.1	12.8
Current	15.5	38.0	37.9
Age at starting to smoke (% of ever smokers)			
30 years	9.6	7.0	5.0†
20-29 years	36.0	28.7	31.9
15-19 years	37.4	41.6	42.9
14 years	17.0	22.7	20.3
Years smoked (% of ever smokers)			
9 years	6.7	4.6	3.3
10-19 years	10.3	5.6	6.0
20-29 years	21.6	14.3	17.0
30-39 years	31.4	29.7	30.8
40 years	30.0	45.8	42.9
Cigarettes smoked per day (% of ever smokers)			
6	16.6	9.1	8.8
7-12	25.4	21.3	19.2
13-22	36.5	39.9	41.8
23-32	9.5	16.1	14.3
33-42	7.2	8.4	8.8
43	4.8	5.2	7.1

*Individuals (N=4,498) with other cough and phlegm symptom outcomes (incident cough or phlegm alone, prevalent cough and/or phlegm, and discordant incident/prevalent combinations) are not included in this table, but are included in the online supplement. See Table E2 for characteristics.

†Percentages do not add up to 100, because of rounding.

‡Chi-square or Students *t*-test *P*-values were <0.01 for all factors comparing the case group to the non-cases, with one exception for age at starting to smoke comparing cases with symptoms for three or more months to non-cases (*P* = 0.056).

§Individuals were categorized as never smokers if they had never smoked at least one cigarette a day for one year or longer.

Table 2 Odds ratios by quartile of daily intake of non-starch polysaccharides, major contributors to non-starch polysaccharide intake and related nutrients, in relation to incident cough with phlegm: The Singapore Chinese Health Study*

	Non-cases N=44,068	All cases of incident cough with phlegm N=571	OR (95% CI)†	Cases with symptoms for 3 months per year N=359	OR (95% CI)†
Total non-starch polysaccharides (g)					
Q1 (4.7)§	9,978	212	1.0 (ref.)	133	1.0 (ref.)
Q2 (6.6)	11,023	154	0.82 (0.66, 1.01)	94	0.81 (0.62, 1.06)
Q3 (8.5)	11,355	106	0.61 (0.48, 0.78)	73	0.69 (0.51, 0.93)
Q4 (11.6)	11,712	99	0.61 (0.47, 0.78)	59	0.60 (0.43, 0.82)
<i>P</i> for trend ‡			<0.001		0.001
<u>Major contributors of non-starch polysaccharides</u>					
Total fruit and fruit juices (g)					
Q1 (44.4)§	10,012	187	1.0 (ref.)	127	1.0 (ref.)
Q2 (127.1)	11,044	131	0.73 (0.58, 0.92)	83	0.70 (0.53, 0.94)
Q3 (210.9)	11,414	128	0.72 (0.56, 0.91)	74	0.64 (0.47, 0.87)
Q4 (377.0)	11,598	125	0.67 (0.52, 0.87)	75	0.64 (0.46, 0.88)
<i>P</i> for trend ‡			0.006		0.010
All grain products (g)					
Q1 (308.1)§	11,249	126	1.0 (ref.)	80	1.0 (ref.)
Q2 (449.7)	11,053	127	0.93 (0.72, 1.21)	88	1.08 (0.78, 1.49)
Q3 (546.8)	11,021	139	0.98 (0.74, 1.29)	79	0.98 (0.69, 1.41)
Q4 (858.0)	10,745	179	1.12 (0.80, 1.56)	112	1.32 (0.87, 2.00)
<i>P</i> for trend ‡			0.301		0.133
Total vegetables and vegetable juices (g)					
Q1 (51.7)§	10,090	151	1.0 (ref.)	95	1.0 (ref.)
Q2 (82.3)	11,087	148	0.98 (0.77, 1.23)	93	1.02 (0.76, 1.37)
Q3 (115.5)	11,358	130	0.87 (0.68, 1.12)	88	1.02 (0.75, 1.39)
Q4 (176.8)	11,533	142	0.92 (0.70, 1.21)	83	0.97 (0.69, 1.37)
<i>P</i> for trend ‡			0.504		0.820
<u>Related nutrients</u>					
Vitamin C (mg)ll					
Q1 (32.1)§	10,011	183	1.0 (ref.)	115	1.0 (ref.)
Q2 (57.2)	10,992	157	0.95 (0.76, 1.18)	95	0.94 (0.72, 1.24)
Q3 (85.7)	11,361	124	0.79 (0.62, 0.99)	82	0.87 (0.65, 1.16)
Q4 (138.6)	11,704	107	0.73 (0.57, 0.94)	67	0.78 (0.57, 1.07)
<i>P</i> for trend ‡			0.006		0.103
Vitamin E (mg)ll					
Q1 (3.9)§	10,118	187	1.0 (ref.)	111	1.0 (ref.)
Q2 (5.0)	10,994	139	0.83 (0.66, 1.04)	83	0.84 (0.63, 1.12)
Q3 (6.0)	11,318	133	0.87 (0.69, 1.09)	97	1.09 (0.82, 1.44)
Q4 (7.5)	11,638	112	0.78 (0.61, 0.99)	68	0.82 (0.60, 1.13)
<i>P</i> for trend ‡			0.063		0.477
Total carotenoids (i g)ll					
Q1 (2,629.4)§	10,033	164	1.0 (ref.)	107	1.0 (ref.)
Q2 (4,098.3)	10,975	141	0.95 (0.75, 1.19)	85	0.90 (0.67, 1.20)
Q3 (5,629.0)	11,409	149	1.07 (0.85, 1.34)	97	1.10 (0.83, 1.46)
Q4 (8,493.8)	11,651	117	0.90 (0.71, 1.16)	70	0.87 (0.64, 1.19)
<i>P</i> for trend ‡			0.567		0.586

Table 2 - continued

Beta-carotene (i g)ll					
Q1 (949.4)§	10,024	165	1.0 (ref.)	105	1.0 (ref.)
Q2 (1,524.1)	11,014	139	0.94 (0.75, 1.18)	91	0.99 (0.74, 1.31)
Q3 (2,140.1)	11,410	141	1.00 (0.80, 1.26)	91	1.05 (0.79, 1.40)
Q4 (3,371.3)	11,620	126	0.98 (0.77, 1.25)	72	0.92 (0.67, 1.25)
<i>P</i> for trend ‡			0.998		0.626

*Individuals (N=4,498) with other symptom outcomes (incident cough or phlegm alone, prevalent cough and/or phlegm, and discordant incident/prevalent combinations) are not included in this table, but their data on the above foods and nutrients can be found in the online supplement (Table E3).

†Models are adjusted for age, total energy intake, dialect group, gender, smoking status (never, former, current), age at starting to smoke (< 20, 15-19, > 14 years), and cigarettes per day (< 12, 13-22, > 23).

These odds ratios are for individual dietary items without adjustment for the other items.

‡Tests for trend were conducted using the median value for each quartile of the specified dietary factor analyzed as an ordinal variable in the regression models.

§Food and energy-adjusted nutrients were categorized into quartiles, based on the distribution of the entire baseline cohort. Median values within quartiles are presented in parentheses, and are based on the subjects included in these analyses.

llNutrient intake from food sources.

Table 3 Odds ratios by tertile of daily intake of specific fruits and incident cough with phlegm: The Singapore Chinese Health Study*

	Non-cases N=44,068	All cases of incident cough with phlegm N=571	OR (95% CI)†	Cases with symptoms for 3 months per year N=359	OR (95% CI)†
Apples (g)					
T1 (0.0)§	13,353	248	1.0 (ref.)	165	1.0 (ref.)
T2 (19.1)	10,796	125	0.75 (0.60, 0.93)	70	0.63 (0.47, 0.84)
T3 (47.8)	19,919	198	0.67 (0.55, 0.82)	124	0.64 (0.50, 0.82)
<i>P</i> for trend‡			<0.001		0.001
Oranges (g)					
T1 (0.0)§	14,019	238	1.0 (ref.)	153	1.0 (ref.)
T2 (21.4)	10,634	121	0.76 (0.61, 0.95)	70	0.69 (0.52, 0.92)
T3 (53.6)	19,415	212	0.79 (0.65, 0.96)	136	0.83 (0.65, 1.06)
<i>P</i> for trend‡			0.026		0.201
Pears (g)					
T1 (0.0)§	16,514	266	1.0 (ref.)	181	1.0 (ref.)
T2 (12.6)	15,057	178	0.84 (0.69, 1.02)	105	0.73 (0.58, 0.94)
T3 (25.2)	12,497	127	0.75 (0.60, 0.93)	73	0.65 (0.49, 0.86)
<i>P</i> for trend‡			0.007		0.001
Grapes (g)					
T1 (0.0)§	23,818	351	1.0 (ref.)	224	1.0 (ref.)
T2 (1.7)	9,703	122	0.94 (0.76, 1.16)	73	0.89 (0.68, 1.17)
T3 (5.8)	10,547	98	0.71 (0.57, 0.90)	62	0.74 (0.56, 0.99)
<i>P</i> for trend‡			0.004		0.043

*Individuals (N=4,498) with other symptom outcomes (incident cough or phlegm, prevalent cough and/or phlegm, and discordant incident/prevalent combinations) are not included in this table, but can be found in the online supplement (Table E4).

†Models are adjusted for age, total energy intake, dialect group, gender, smoking status (never, former, current), age at starting to smoke (< 20, 15-19, ≥ 14 years), and cigarettes per day (< 12, 13-22, ≥ 23).

‡Tests for trend were conducted using the median value for each tertile of the specified dietary factor analyzed as an ordinal variable in the regression models.

§Fruits were categorized into tertiles, based on the distribution of the entire baseline cohort. Median values within tertiles are presented in parentheses, and are based on the subjects included in these analyses. Tertiles rather than quartiles were used because consumption of any individual fruit is substantially less than total fruits.

Table 4 Odds ratios by quartile of daily intake of soyfoods and soy isoflavones in relation to incident cough with phlegm: The Singapore Chinese Health Study*

	Non-cases N=44,068	All cases of incident cough with phlegm N=571	OR (95% CI)†	Cases with symptoms for 3 months per year N=359	OR (95% CI)†
Total soyfoods (g)‡					
Q1 (31.4)§	10,457	163	1.0 (ref.)	102	1.0 (ref.)
Q2 (67.9)	11,080	147	0.89 (0.71, 1.12)	96	0.95 (0.71, 1.26)
Q3 (112.0)	11,284	136	0.80 (0.63, 1.02)	81	0.80 (0.59, 1.09)
Q4 (205.7)	11,247	125	0.69 (0.53, 0.90)	80	0.76 (0.55, 1.06)
<i>P</i> for trend			0.005		0.085
Total soy isoflavones (mg)					
Q1 (5.4)§	10,483	183	1.0 (ref.)	112	1.0 (ref.)
Q2 (11.4)	10,987	156	0.91 (0.73, 1.13)	99	0.95 (0.72, 1.24)
Q3 (17.9)	11,188	126	0.76 (0.61, 0.96)	78	0.78 (0.58, 1.04)
Q4 (30.3)	11,410	106	0.67 (0.53, 0.86)	70	0.73 (0.54, 0.98)
<i>P</i> for trend			0.001		0.022

*Individuals (N=4,498) with other symptom outcomes (incident cough or phlegm, prevalent cough and/or phlegm, and discordant incident/prevalent combinations) are not included in this table, but can be found in the online supplement (Table E5).

†Models are adjusted for age, total energy intake, dialect group, gender, smoking status (never, former, current), age at starting to smoke (< 20, 15-19, ≥ 14 years), and cigarettes per day (< 12, 13-22, ≥ 23). These odds ratios are for individual dietary items without adjustment for the other item.

‡Expressed in units of tofu-equivalent (see “Methods” section for details).

§Food and energy-adjusted nutrients were categorized into quartiles, based on the distribution of the entire baseline cohort. Median values within quartiles are presented in parentheses, and are based on the subjects included in these analyses.

||Tests for trend were conducted using the median value for each quartile of the specified dietary factor analyzed as an ordinal variable in the regression models.

Table 5 Odds ratios for daily intake of selected dietary factors and incident cough with phlegm, with and without adjustment for total non-starch polysaccharides: The Singapore Chinese Health Study*

	OR (95% CI) for dietary factor† Adjusted for covariates only‡		OR (95% CI) for dietary factor† Adjusted for covariates‡ and non- starch polysaccharides	
	All cases of incident cough with phlegm	Cases with symptoms for 3 months per year	All cases of incident cough with phlegm§	Cases with symptoms for 3 months per year§
<u>Dietary factors</u>				
Total fruit	0.67 (0.52, 0.87)	0.64 (0.46, 0.88)	1.06 (0.74, 1.52)	0.89 (0.57, 1.39)
Apples	0.67 (0.55, 0.82)	0.64 (0.50, 0.82)	0.79 (0.64, 0.98)	0.74 (0.56, 0.97)
Pears	0.75 (0.60, 0.93)	0.65 (0.49, 0.86)	0.86 (0.69, 1.09)	0.73 (0.54, 0.98)
Grapes	0.71 (0.57, 0.90)	0.74 (0.56, 0.99)	0.79 (0.62, 0.99)	0.81 (0.61, 1.09)
All grain products	1.12 (0.80, 1.56)	1.32 (0.87, 2.00)	0.89 (0.63, 1.26)	1.08 (0.70, 1.67)
Total vegetables	0.92 (0.70, 1.21)	0.97 (0.69, 1.37)	1.23 (0.91, 1.65)	1.27 (0.87, 1.85)
Vitamin C II	0.73 (0.57, 0.94)	0.78 (0.57, 1.07)	1.16 (0.82, 1.65)	1.30 (0.84, 2.02)
Vitamin E II	0.78 (0.61, 0.99)	0.82 (0.60, 1.13)	1.08 (0.81, 1.43)	1.16 (0.81, 1.67)
Total soyfoods	0.69 (0.53, 0.90)	0.76 (0.55, 1.06)	0.83 (0.62, 1.10)	0.91 (0.64, 1.30)
Total soy isoflavones	0.67 (0.53, 0.86)	0.73 (0.54, 0.98)	0.80 (0.62, 1.04)	0.86 (0.62, 1.19)
Non-starch polysaccharides	0.61 (0.47, 0.78)	0.60 (0.43, 0.82)	---	---

*Individuals (N=4,498) with other symptom outcomes (incident cough or phlegm, prevalent cough and/or phlegm, and discordant incident/prevalent combinations) are not included in this table, but can be found in the online supplement (Table E6).

†Odds ratios (OR) and 95% confidence intervals (CI) for fourth versus first quartile of daily intake for all specified dietary factors, except for apples, pears and grapes, where ORs and CIs are for third versus first tertile of daily intake.

‡Covariates include age, total energy intake, dialect group, gender, smoking status (never, former, current), age at starting to smoke (< 20, 15-19, ≥ 14 years), and cigarettes per day (< 12, 13-22, ≥ 23).

§Odds ratios for non-starch polysaccharide intake in each model were statistically significant and ranged from 0.49 to 0.67.

|||Nutrient intake from food sources.

ON-LINE SUPPLEMENT DATA

Dietary fiber and reduced cough with phlegm: a cohort study in Singapore

Lesley M. Butler, Woon-Puay Koh, Hin-Peng Lee, Mimi C. Yu, Stephanie J. London

Sections from the Singapore Chinese Health Study Questionnaire

SMOKING

1. Have you ever smoked at least one cigarette a day for 1 year or longer?
2. At what age did you start smoking cigarettes on a regular basis?
3. What is the total number of years that you smoked cigarettes on a regular basis?
4. If you quit smoking cigarettes, how long ago did you quit?
5. If you smoked manufactured cigarettes, what is the average number of cigarettes that you smoked per day?
6. If you smoked self-rolled cigarettes, how many tahils/liangs of tobacco, on average, did you smoke in a month?

HISTORY OF RESPIRATORY DISEASES (E1)

COUGH

1. Do you usually cough when you get up in the morning? (Exclude clearing of the throat)
2. Do you usually cough at all during the rest of the day or at night?
3. Do you usually cough for 3 months or more during the year?
4. For how many years have you had this cough?

SPUTUM / PHLEGM

1. Do you usually bring up phlegm from your chest when you get up in the morning?
(Exclude phlegm from the nose, count phlegm swallowed)
2. Do you usually bring up phlegm from your chest during the rest of the day or at night?
3. Do you usually bring up phlegm from your chest on most days for three months or more during the year?
4. For how many years have you had trouble with phlegm?

Table E1 Characteristics of the Singapore Chinese Health Study, among participants with and without respiratory follow-up data.

	With respiratory follow-up N=49,140	Without respiratory follow-up N=14,117*
<u>Demographics</u>		
Sex (%)		
Men	42.3	50.7
Women	57.7	49.3
Age		
Mean years (standard deviation)	55.8 (7.7)	58.9 (8.5)
Dialect group (%)		
Hokkien	50.6	64.6
Cantonese	49.4	35.4
<u>Smoking History</u>		
Smoking status (%)		
Never†	72.2‡	60.0
Past	10.5	13.1
Current	17.4	26.9
Age at starting to smoke (% of ever smokers)		
30 years	9.2‡	9.2‡
20-29 years	35.7	32.0
15-19 years	37.7	38.8
14 years	17.5	20.1
Years smoked (% of ever smokers)		
9 years	6.3‡	3.7‡
10-19 years	9.7	7.1
20-29 years	21.4	16.6
30-39 years	31.4	28.3
40 years	31.3	44.4
Cigarettes smoked per day (% of ever smokers)		
6	15.8‡	14.9
7-12	25.0	22.9
13-22	37.4	39.4
23-32	9.9	9.6
33-42	7.4	7.8
43	4.6	5.4
<u>Mean daily dietary intake (standard deviation)§</u>		
Total non-starch polysaccharide (g)	8.5 (2.8)	7.9 (3.0)
Total fruit (g)	207.2 (72.5)	185.8 (78.3)
Total vegetables (g)	112.4 (33.1)	104.0 (35.9)
Vitamin C (mg)¶	88.0 (29.2)	79.4 (31.3)
Soy foods (g)¶¶	113.4 (44.0)	105.4 (47.5)
Total soy isoflavones (mg)	19.0 (7.4)	17.6 (8.0)

* Includes 4,871 subjects who died prior to the follow-up interview.

† Individuals were categorized as never smokers if they had never smoked at least one cigarette a day for one year or longer.

‡ Percentages do not add up to 100, because of rounding.

§ Foods and nutrients are adjusted for age, dialect group, gender, smoking status and total calories.

¶ Nutrient intake from food sources.

¶¶ Expressed in units of tofu-equivalent (see “Methods” section for details).

Table E2 Characteristics of the Singapore Chinese Health Study according to the presence of cough and/or phlegm*

	Non-cases	Incident cough only	Incident phlegm only	Prevalent cough with phlegm	Miscellaneous cough and/or phlegm
	N=44,068	N=512	N=1,271	N=668	N=2,047
Sex (%)					
Men	40.3	50.8	58.2	66.2	61.1
Women	59.7	49.2	41.8	33.8	38.9
Chi-square <i>P</i> -value†		<0.0001	<0.0001	<0.0001	<0.0001
Age					
Mean years (standard deviation)	55.7 (7.7)	56.8 (7.7)	57.1 (8.1)	57.4 (8.3)	55.7 (7.8)
Students t-test <i>P</i> -value†		0.001	<0.0001	<0.0001	0.883
Dialect group (%)					
Hokkien	51.1	50.8	44.7	48.1	41.4
Cantonese	48.9	49.2	55.3	52.0	58.6
Chi-square <i>P</i> -value†		0.884	<0.0001	0.117	<0.0001
Smoking status (%)					
Never‡	74.3	64.8	54.3	42.8	55.2
Past	10.2	9.2	11.8	13.5	13.6
Current	15.5	26.0	33.9	43.7	31.2
Chi-square <i>P</i> -value†		<0.0001	<0.0001	<0.0001	<0.0001
Age at starting to smoke (% of ever smokers)					
30 years	9.6	7.8	7.9	6.0	6.4
20-29 years	36.0	36.7	41.0	31.4	32.6
15-19 years	37.4	37.8	32.9	38.5	41.8
14 years	17.0	17.8	18.2	24.1	19.2
Chi-square <i>P</i> -value†		0.864	0.027	0.0004	0.0003
Years smoked (% of ever smokers)					
9 years	6.7	5.6	4.3	2.9	4.4
10-19 years	10.3	6.1	6.9	5.8	7.3
20-29 years	21.6	20.6	21.7	19.4	21.2
30-39 years	31.4	33.3	30.8	30.1	33.2
40 years	30.0	34.4	36.3	41.9	34.0
Chi-square <i>P</i> -value†		0.305	0.001	<0.0001	0.0005
Cigarettes smoked per day (% of ever smokers)					
6	16.6	9.4	12.9	10.2	12.1
7-12	25.4	29.4	23.6	19.9	22.9
13-22	36.5	40.0	42.9	47.4	39.9
23-32	9.5	11.7	10.0	12.8	11.2
33-42	7.2	6.7	7.2	7.3	8.7

43	4.8	2.8	3.4	2.4	5.1
Chi-square <i>P</i> -value†		0.084	0.017	<0.0001	0.0008

*Individuals (N=571) with incident cough plus phlegm are not included in this table, but can be found in the manuscript (Table 1).

†Chi-square or Students *t*-test *P*-values were calculated for all factors comparing each case group to the non-cases.

‡Individuals were categorized as never smokers if they had never smoked at least one cigarette a day for one year or longer.

Table E3 Odds ratios by quartile of daily intake of non-starch polysaccharides, major contributors to non-starch polysaccharide intake and related nutrients, in relation to cough and/or phlegm: The Singapore Chinese Health Study*

	Non-cases N=44,068	Incident cough only N=512	OR (95% CI)†	Incident phlegm only N=1,271	OR (95% CI)†	Prevalent cough with phlegm N=668	OR (95% CI)†	Miscellaneous cough and/or phlegm N=2,047	OR (95% CI)†
Total non-starch polysaccharides (g)									
Q1 (4.7)§	9,978	120	1.0 (ref.)	343	1.0 (ref.)	239	1.0 (ref.)	557	1.0 (ref.)
Q2 (6.6)	11,023	134	1.13 (0.88, 1.45)	328	1.04 (0.89, 1.22)	174	0.86 (0.70, 1.05)	526	1.00 (0.88, 1.13)
Q3 (8.5)	11,355	132	1.14 (0.89, 1.47)	315	1.07 (0.91, 1.25)	156	0.85 (0.69, 1.05)	517	1.03 (0.91, 1.17)
Q4 (11.6)	11,712	126	1.11 (0.85, 1.44)	285	1.01 (0.85, 1.19)	99	0.59 (0.46, 0.75)	447	0.93 (0.81, 1.06)
<i>P</i> for trend ‡			0.532		0.954		<0.001		0.311
Major contributors of non-starch polysaccharides									
Total fruit and fruit juices (g)									
Q1 (44.4)§	10,012	121	1.0 (ref.)	324	1.0 (ref.)	209	1.0 (ref.)	471	1.0 (ref.)
Q2 (127.1)	11,044	129	1.02 (0.79, 1.32)	314	0.95 (0.81, 1.12)	161	0.77 (0.62, 0.95)	501	0.95 (0.83, 1.08)
Q3 (210.9)	11,414	128	0.99 (0.76, 1.28)	300	0.89 (0.75, 1.05)	163	0.76 (0.61, 0.94)	512	0.90 (0.79, 1.03)
Q4 (377.0)	11,598	134	1.01 (0.77, 1.32)	333	0.97 (0.81, 1.15)	135	0.59 (0.46, 0.75)	563	0.90 (0.79, 1.04)
<i>P</i> for trend ‡			0.991		0.780		<0.001		0.173
All grain products (g)									
Q1 (308.1)§	11,249	123	1.0 (ref.)	263	1.0 (ref.)	112	1.0 (ref.)	415	1.0 (ref.)
Q2 (449.7)	11,053	108	0.82 (0.63, 1.08)	300	1.05 (0.88, 1.25)	175	1.33 (1.03, 1.71)	450	0.90 (0.78, 1.04)
Q3 (546.8)	11,021	140	1.02 (0.77, 1.35)	342	1.12 (0.92, 1.35)	172	1.16 (0.88, 1.52)	527	0.89 (0.76, 1.03)
Q4 (858.0)	10,745	141	0.94 (0.66, 1.34)	366	1.06 (0.84, 1.33)	209	1.13 (0.82, 1.56)	655	0.84 (0.70, 1.01)
<i>P</i> for trend ‡			0.960		0.803		0.946		0.107
Total vegetables and vegetable juices (g)									
Q1 (51.7)§	10,090	120	1.0 (ref.)	319	1.0 (ref.)	165	1.0 (ref.)	428	1.0 (ref.)
Q2 (82.3)	11,087	137	1.05 (0.82, 1.36)	326	0.97 (0.83, 1.14)	175	1.01 (0.81, 1.26)	489	1.01 (0.88, 1.16)
Q3 (115.5)	11,358	141	1.05 (0.81, 1.37)	325	0.95 (0.81, 1.13)	179	1.03 (0.82, 1.29)	526	1.03 (0.90, 1.18)
Q4 (176.8)	11,533	114	0.81 (0.60, 1.09)	301	0.85 (0.70, 1.02)	149	0.79 (0.61, 1.02)	604	1.07 (0.93, 1.24)
<i>P</i> for trend ‡			0.099		0.065		0.051		0.308
Related nutrients									
Vitamin C (mg)¶									
Q1 (32.1)§	10,011	131	1.0 (ref.)	320	1.0 (ref.)	235	1.0 (ref.)	529	1.0 (ref.)
Q2 (57.2)	10,992	121	0.91 (0.71, 1.17)	311	0.99 (0.84, 1.16)	168	0.77 (0.63, 0.94)	529	0.96 (0.85, 1.09)
Q3 (85.7)	11,361	129	0.97 (0.75, 1.24)	323	1.04 (0.89, 1.22)	162	0.77 (0.63, 0.95)	502	0.91 (0.80, 1.03)
Q4 (138.6)	11,704	131	1.00 (0.78, 1.28)	317	1.06 (0.90, 1.25)	103	0.54 (0.42, 0.68)	487	0.92 (0.81, 1.05)
<i>P</i> for trend ‡			0.803		0.362		<0.001		0.207

Vitamin E (mg)II									
Q1 (3.9)§	10,118	137	1.0 (ref.)	353	1.0 (ref.)	203	1.0 (ref.)	534	1.0 (ref.)
Q2 (5.0)	10,994	128	0.95 (0.74, 1.21)	323	1.00 (0.85, 1.16)	189	1.09 (0.89, 1.34)	514	1.05 (0.92, 1.19)
Q3 (6.0)	11,318	125	0.95 (0.74, 1.22)	305	1.01 (0.86, 1.18)	151	0.98 (0.79, 1.21)	480	1.03 (0.91, 1.18)
Q4 (7.5)	11,638	122	0.95 (0.73, 1.22)	290	1.01 (0.86, 1.19)	125	0.88 (0.70, 1.12)	519	1.16 (1.01, 1.32)
<i>P</i> for trend ‡			0.707		0.859		0.224		0.037
Total carotenoids (i g)II									
Q1 (2,629.4)§	10,033	133	1.0 (ref.)	339	1.0 (ref.)	206	1.0 (ref.)	521	1.0 (ref.)
Q2 (4,098.3)	10,975	118	0.88 (0.68, 1.13)	345	1.05 (0.90, 1.23)	178	0.95 (0.78, 1.17)	510	0.98 (0.86, 1.11)
Q3 (5,629.0)	11,409	133	1.00 (0.78, 1.27)	312	0.98 (0.83, 1.15)	153	0.88 (0.71, 1.09)	505	0.98 (0.87, 1.12)
Q4 (8,493.8)	11,651	128	0.98 (0.77, 1.26)	275	0.90 (0.76, 1.06)	131	0.81 (0.65, 1.02)	511	1.03 (0.91, 1.17)
<i>P</i> for trend ‡			0.826		0.124		0.058		0.573
Beta-carotene (i g)II									
Q1 (949.4)§	10,024	128	1.0 (ref.)	342	1.0 (ref.)	195	1.0 (ref.)	532	1.0 (ref.)
Q2 (1,524.1)	11,014	129	1.00 (0.78, 1.29)	334	1.03 (0.88, 1.20)	192	1.11 (0.91, 1.36)	509	0.98 (0.87, 1.11)
Q3 (2,140.1)	11,410	137	1.07 (0.84, 1.37)	324	1.02 (0.87, 1.20)	156	0.96 (0.77, 1.20)	505	0.99 (0.87, 1.13)
Q4 (3,371.3)	11,620	118	0.95 (0.74, 1.23)	271	0.91 (0.77, 1.08)	125	0.86 (0.68, 1.09)	501	1.05 (0.92, 1.19)
<i>P</i> for trend ‡			0.731		0.228		0.098		0.407

*Individuals (N=571) with incident cough plus phlegm are not included in this table, but their data on the above foods and nutrients can be found in the manuscript (Table 2).

†Models are adjusted for age, total energy intake, dialect group, gender, smoking status (never, former, current), age at starting to smoke (20, 15-19, 14 years), and cigarettes per day (12, 13-22, 23). These odds ratios are for individual dietary items without adjustment for the other items.

‡Tests for trend were conducted using the median value for each quartile of the specified dietary factor analyzed as an ordinal variable in the regression models.

§Food and energy-adjusted nutrients were categorized into quartiles, based on the distribution of the entire baseline cohort. Median values within quartiles are presented in parentheses, and are based on subjects included in these analyses.

IIINutrient intake from food sources.

Table E4 Odds ratios by tertile of daily intake of specific fruits in relation to cough and/or phlegm: The Singapore Chinese Health Study*

	Non-cases N=44,068	Incident cough only N=512	OR (95% CI)†	Incident phlegm only N=1,271	OR (95% CI)†	All prevalent cough with phlegm N=668	OR (95% CI)†	Miscellaneous cough and/or phlegm N=2,047	OR (95% CI)†
Apples (g)									
T1 (0.0)‡	13,353	191	1.0 (ref.)	453	1.0 (ref.)	287	1.0 (ref.)	689	1.0 (ref.)
T2 (19.1)	10,796	126	0.88 (0.70, 1.10)	314	0.98 (0.85, 1.14)	149	0.79 (0.64, 0.96)	503	0.99 (0.88, 1.11)
T3 (47.8)	19,919	195	0.74 (0.60, 0.91)	504	0.90 (0.79, 1.03)	232	0.71 (0.59, 0.85)	855	0.93 (0.83, 1.03)
<i>P</i> for trend§			0.005		0.124		<0.001		0.159
Oranges (g)									
T1 (0.0)‡	14,019	188	1.0 (ref.)	441	1.0 (ref.)	317	1.0 (ref.)	729	1.0 (ref.)
T2 (21.4)	10,634	128	0.94 (0.74, 1.17)	314	1.00 (0.86, 1.16)	133	0.61 (0.49, 0.75)	491	0.90 (0.80, 1.01)
T3 (53.6)	19,415	196	0.79 (0.64, 0.98)	516	0.94 (0.82, 1.07)	218	0.57 (0.48, 0.69)	827	0.81 (0.73, 0.90)
<i>P</i> for trend§			0.026		0.306		<0.001		<0.001
Pears (g)									
T1 (0.0)‡	16,514	207	1.0 (ref.)	489	1.0 (ref.)	315	1.0 (ref.)	793	1.0 (ref.)
T2 (12.6)	15,057	166	0.93 (0.76, 1.15)	443	1.11 (0.97, 1.26)	199	0.81 (0.67, 0.97)	679	1.01 (0.91, 1.12)
T3 (25.2)	12,497	139	0.95 (0.76, 1.19)	339	1.05 (0.91, 1.22)	154	0.77 (0.63, 0.94)	575	1.00 (0.89, 1.12)
<i>P</i> for trend§			0.619		0.412		0.005		0.977
Grapes (g)									
T1 (0.0)‡	23,818	297	1.0 (ref.)	706	1.0 (ref.)	407	1.0 (ref.)	1,125	1.0 (ref.)
T2 (1.7)	9,703	96	0.83 (0.65, 1.04)	247	0.92 (0.79, 1.07)	127	0.85 (0.69, 1.04)	444	1.01 (0.91, 1.14)
T3 (5.8)	10,547	119	0.95 (0.76, 1.18)	318	1.11 (0.97, 1.27)	134	0.84 (0.68, 1.02)	478	0.97 (0.86, 1.08)
<i>P</i> for trend§			0.626		0.140		0.076		0.570

*Individuals (N=571) with incident cough plus phlegm are not included in this table, but their data for individual fruits can be found in the manuscript (Table 3).

†Models are adjusted for age, total energy intake, dialect group, gender, smoking status (never, former, current), age at starting to smoke (20, 15-19, 14 years), and cigarettes per day (12, 13-22, 23). These odds ratios are for individual fruit items without adjustment for the other items.

‡Fruits were categorized into tertiles, based on the distribution of the entire baseline cohort. Median values within tertiles are presented in parentheses, and are based on the subset of the cohort used in these analyses. Tertiles rather than quartiles were used because consumption of any individual fruit is substantially less than total fruits.

§Tests for trend were conducted using the median value for each tertile of the specified dietary factor analyzed as an ordinal variable in the regression models.

Table E5 Odds ratios by quartile of daily intake of soyfoods and soy isoflavones in relation to cough and/or phlegm: The Singapore Chinese Health Study*

	Non-cases N=44,068	Incident cough only N=512	OR (95% CI)†	Incident phlegm only N=1,271	OR (95% CI)†	All prevalent cough with phlegm N=668	OR (95% CI)†	Miscellaneous cough and/or phlegm N=2,047	OR (95% CI)†
Total soyfoods (g)‡									
Q1 (31.4)§	10,457	116	1.0 (ref.)	308	1.0 (ref.)	178	1.0 (ref.)	460	1.0 (ref.)
Q2 (67.9)	11,080	117	0.98 (0.76, 1.28)	305	0.98 (0.84, 1.16)	170	0.94 (0.76, 1.16)	494	0.99 (0.87, 1.13)
Q3 (112.0)	11,284	144	1.20 (0.93, 1.56)	327	1.05 (0.89, 1.24)	155	0.83 (0.66, 1.04)	525	0.99 (0.87, 1.13)
Q4 (205.7)	11,247	135	1.12 (0.84, 1.48)	331	1.06 (0.88, 1.26)	165	0.82 (0.65, 1.05)	568	0.98 (0.85, 1.13)
<i>P</i> for trend			0.341		0.435		0.104		0.832
Total soy isoflavones (mg)									
Q1 (5.4)§	10,483	121	1.0 (ref.)	343	1.0 (ref.)	206	1.0 (ref.)	568	1.0 (ref.)
Q2 (11.4)	10,987	119	0.99 (0.77, 1.28)	290	0.89 (0.76, 1.04)	180	0.95 (0.77, 1.16)	522	0.94 (0.83, 1.07)
Q3 (17.9)	11,188	136	1.15 (0.90, 1.48)	320	1.02 (0.88, 1.20)	153	0.86 (0.70, 1.07)	499	0.94 (0.83, 1.07)
Q4 (30.3)	11,410	136	1.18 (0.92, 1.51)	318	1.07 (0.91, 1.25)	129	0.79 (0.63, 0.99)	458	0.91 (0.80, 1.03)
<i>P</i> for trend			0.124		0.154		0.027		0.172

*Individuals (N=571) with incident cough plus phlegm are not included in this table, but their data for soyfoods and soy isoflavones can be found in the manuscript (Table 4).

†Models are adjusted for age, total energy intake, dialect group, gender, smoking status (never, former, current), age at starting to smoke (< 20, 15-19, 14 years), and cigarettes per day (< 12, 13-22, ≥ 23). These odds ratios are for individual dietary items without adjustment for the other items.

‡Expressed in units of tofu-equivalent (see “Methods” section for details).

§Food and energy-adjusted nutrients were categorized into quartiles, based on the distribution of the entire baseline cohort. Median values within quartiles are presented in parentheses, and are based on the subjects included in these analyses.

||Tests for trend were conducted using the median value for each quartile of the specified dietary factor analyzed as an ordinal variable in the regression models.

Table E6 Odds ratios for daily intake of selected dietary factors in relation to cough and/or phlegm, with and without adjustment for total non-starch polysaccharides: The Singapore Chinese Health Study*

	OR (95% CI) for dietary factor† Adjusted for covariates only‡				OR (95% CI) for dietary factor† Adjusted for covariates‡ and non-starch polysaccharides			
	Incident cough only	Incident phlegm only	Prevalent cough with phlegm	Miscellaneous cough and/or phlegm	Incident cough only§	Incident phlegm only	Prevalent cough with phlegm¶	Miscellaneous cough and/or phlegm**
Dietary factors								
Total fruit	1.01(0.77, 1.32)	0.97 (0.81, 1.15)	0.59 (0.46, 0.75)	0.90 (0.79, 1.04)	0.87 (0.59, 1.27)	0.90 (0.71, 1.15)	0.70 (0.50, 0.97)	0.88 (0.73, 1.07)
Apples	0.74 (0.60, 0.91)	0.90 (0.79, 1.03)	0.71 (0.59, 0.85)	0.93 (0.83, 1.03)	0.67 (0.54, 0.84)	0.88 (0.76, 1.01)	0.79 (0.65, 0.96)	0.93 (0.83, 1.05)
Pears	0.95 (0.76, 1.19)	1.05 (0.91, 1.22)	0.77 (0.63, 0.94)	1.00 (0.89, 1.12)	0.92 (0.73, 1.16)	1.05 (0.90, 1.22)	0.85 (0.69, 1.05)	1.01 (0.90, 1.14)
Grapes	0.95 (0.76, 1.18)	1.11 (0.97, 1.27)	0.84 (0.68, 1.02)	0.97 (0.86, 1.08)	0.92 (0.74, 1.16)	1.11 (0.96, 1.28)	0.90 (0.73, 1.10)	0.98 (0.87, 1.09)
All grain products	0.94 (0.66, 1.34)	1.06 (0.84, 1.33)	1.13 (0.82, 1.56)	0.84 (0.70, 1.01)	0.98 (0.68, 1.41)	1.07 (0.84, 1.36)	0.92 (0.66, 1.29)	0.80 (0.66, 0.97)
Total vegetables	0.81 (0.60, 1.09)	0.85 (0.70, 1.02)	0.79 (0.61, 1.02)	1.07 (0.93, 1.24)	0.73 (0.53, 1.01)	0.81 (0.66, 0.99)	0.96 (0.72, 1.26)	1.12 (0.95, 1.31)
Vitamin C ††	1.00 (0.78, 1.28)	1.06 (0.90, 1.25)	0.54 (0.42, 0.68)	0.92 (0.81, 1.05)	0.89 (0.62, 1.26)	1.12 (0.89, 1.41)	0.58 (0.42, 0.81)	0.91 (0.76, 1.09)
Vitamin E ††	0.95 (0.73, 1.22)	1.01 (0.86, 1.19)	0.88 (0.70, 1.12)	1.16 (1.01, 1.32)	0.86 (0.64, 1.16)	1.00 (0.83, 1.22)	1.17 (0.89, 1.53)	1.26 (1.08, 1.47)
Total soyfoods	1.12 (0.84, 1.48)	1.06 (0.88, 1.26)	0.82 (0.65, 1.05)	0.98 (0.85, 1.13)	1.09 (0.81, 1.46)	1.05 (0.87, 1.27)	0.95 (0.74, 1.23)	1.00 (0.86, 1.16)
Total soy isoflavones	1.18 (0.92, 1.51)	1.07 (0.91, 1.25)	0.79 (0.63, 0.99)	0.91 (0.80, 1.03)	1.16 (0.89, 1.51)	1.08 (0.91, 1.28)	0.91 (0.72, 1.16)	0.91 (0.79, 1.05)
Non-starch polysaccharides	1.11 (0.85, 1.44)	1.01 (0.85, 1.19)	0.59 (0.49, 0.75)	0.93 (0.81, 1.06)	---	---	---	---

*Individuals (N=571) with incident cough plus phlegm are not included in this table, but their data can be found in the manuscript (Table 5).

†Odds ratios (OR) and 95% confidence intervals (CI) for fourth versus first quartile of daily intake for all specified dietary factors, except for apples, pears and grapes, where ORs and CIs are for third versus first tertile of daily intake. Tertiles rather than quartiles were used because consumption of any individual fruit is substantially less than total fruits.

‡Covariates include age, total energy intake, dialect group, gender, smoking status (never, former, current), age at starting to smoke (< 20, 15-19, 14 years), and cigarettes per day (< 12, 13-22, > 23).

§Odds ratios for non-starch polysaccharide intake in each model were not statistically significant and ranged from 1.04 to 1.33.

||Odds ratios for non-starch polysaccharide intake in each model were not statistically significant and ranged from 0.92 to 1.08.

¶Odds ratios for non-starch polysaccharide (NSP) intake in each model were statistically significant and ranged from 0.55 to 0.84, with exceptions for the models with total fruit (OR=0.73; 95% CI:0.52, 1.01; p-value=0.059) and with vitamin C (OR=0.84; 95% CI: 0.60, 1.17; p-value=0.307).

**Odds ratios for non-starch polysaccharide intake in each model were not statistically significant and ranged from 0.83 to 1.03, with one exception for the model with vitamin E (OR=0.83; 95% CI: 0.71, 0.97; p-value=0.017).

††Nutrient intake from food sources.

Table E7 Asthma outcomes by incident cough with phlegm case status: The Singapore Chinese Health Study

	Non-cases N=44,068	All cases of incident cough with phlegm N=571	Cases with symptoms for 3 months per year N=359
Asthma outcomes			
Incident (%)	0.6	5.6	7.2
Prevalent (%)	3.3	6.3	6.4
Never (%)	96.1	88.1	86.4

On-line supplement references

- E1. American Thoracic Society. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1995;152:S78-121.